

## SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR XTENDIMAX™ WITH VAPORGRIP™ TECHNOLOGY BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using XtendiMax™ With VaporGrip™ Technology as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the container label and booklet provided with the product container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.

# XTENDIMAX™

With VaporGrip™  
Technology

This supplemental label expires on 11/09/2018 and must not be used or distributed after this date.

EPA Reg. No. 524-617

GROUP	4	HERBICIDE
-------	---	-----------

## FOR PREEMERGENCE AND POSTEMERGENCE USE ON COTTON WITH XTENDFLEX® TECHNOLOGY

Keep out of reach of children  
**CAUTION!**

In case of an emergency involving this product, call collect, day or night, 314-694-4000.

Bollgard II®, Bollgard® 3, Roundup Ready®, Roundup Ready 2 Xtend®, XtendiMax™, XtendFlex®, and VaporGrip™ are trademarks of Monsanto Technology LLC. All other trademarks are the property of their respective owners.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of herbicide application.

COTTON WITH XTENDFLEX® TECHNOLOGY (INCLUDING BOLLGARD II® XTENDFLEX® COTTON, BOLLGARD® 3 XTENDFLEX® COTTON, OR XTENDFLEX® COTTON) CONTAINS A PATENTED GENE THAT PROVIDES TOLERANCE TO DICAMBA, THE ACTIVE INGREDIENT IN THIS PRODUCT. THIS PRODUCT WILL CAUSE SEVERE CROP INJURY OR DESTRUCTION AND YIELD LOSS IF APPLIED TO COTTON THAT IS NOT DICAMBA TOLERANT, INCLUDING COTTON WITH A TRAIT ENGINEERED TO CONFER TOLERANCE TO AUXIN HERBICIDES OTHER THAN DICAMBA. FOLLOW THE REQUIREMENTS SET FORTH HEREIN TO PREVENT SEVERE CROP INJURY OR DESTRUCTION AND YIELD LOSS. CONTACT WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, OR ANY DESIRABLE PLANTS THAT DO NOT CONTAIN A DICAMBA TOLERANCE GENE OR ARE NOT NATURALLY TOLERANT TO DICAMBA, COULD RESULT IN SEVERE PLANT INJURY OR DESTRUCTION.

Information on cotton with XtendFlex® technology can be obtained from your seed supplier or Monsanto representative. Cotton seed with XtendFlex® technology must be purchased from an authorized licensed seed supplier.

The instructions contained in this Monsanto Supplemental Label include all applications of XtendiMax™ With VaporGrip™ Technology that may be made to cotton with XtendFlex® technology during the cropping season. DO NOT combine these instructions with other instructions in the "COTTON" Section of any other XtendiMax™ With VaporGrip™ Technology label for use over crops that do not contain the dicamba tolerance trait.

Note: Cotton with XtendFlex® technology and methods of controlling weeds and applying dicamba in a cotton with XtendFlex® technology crop are protected under U.S. patent law. A license to use cotton seed with XtendFlex® technology must be obtained prior to use. No license to use cotton with XtendFlex® technology is granted or implied with the purchase of this herbicide product. Cotton with XtendFlex® technology is owned by Monsanto and a license must be obtained from Monsanto before using it. Contact your Authorized Monsanto Retailer for information on obtaining a license to cotton with XtendFlex® technology.

See the "PRODUCT INFORMATION" and "APPLICATION EQUIPMENT AND TECHNIQUES" sections of the XtendiMax™ With VaporGrip™ Technology product label for important use information. In the event that there are any inconsistencies with the directions for use between this supplemental label and any other labeling for this product, follow the directions for use on this supplemental label.

Training and education on proper pesticide application is encouraged. Applicators should visit [www.xtendimaxapplicationrequirements.com](http://www.xtendimaxapplicationrequirements.com) for training information and opportunities relevant to this product.

**TYPES OF APPLICATIONS:** Preplant; At-Planting; Preemergence; Postemergence (In-crop)

XtendiMax™ With VaporGrip™ Technology is approved by U.S. EPA to be used in the following states, subject to county restriction as noted: Alabama, Arkansas, Arizona, Colorado, Delaware, Florida (excluding Palm Beach County), Georgia, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maryland, Michigan, Minnesota, Mississippi, Missouri, Nebraska, New Jersey, New Mexico, New York, North Carolina, North Dakota, Oklahoma, Ohio, Pennsylvania, South Carolina, South Dakota, Tennessee (excluding Wilson County), Texas, Virginia, West Virginia, Wisconsin.

### Restrictions

- Do not apply this product aerially.
- Do not make application of this product if rain is expected in the next 24 hours.

### USE INSTRUCTIONS

Apply this product in a minimum of 10 gallons of spray solution per acre as a broadcast application. For best performance, control weeds early when they are less than 4 inches. Timely application will improve control and reduce weed competition. Refer to the following table for maximum application rates of this product with cotton with XtendFlex® technology.

Maximum Application Rates	
Combined total per year for all applications	88 fluid ounces per acre (2.0 lb. a.e. dicamba per acre)
Total of all Preplant, At-Planting, and Preemergence applications	44 fluid ounces per acre (1.0 lb. a.e. dicamba per acre)
Total of all In-crop applications from emergence up to 7 days pre-harvest	88 fluid ounces per acre (2.0 lb. a.e. dicamba per acre)
Maximum In-crop, single application	22 fluid ounces per acre (0.5 lb. a.e. dicamba per acre)

a.e. – acid equivalent

Refer to Table 1 of the XtendiMax™ With VaporGrip™ Technology label booklet for application rates for weed type and growth stage controlled by this product. Maximum in-crop application rate should be used when treating tough to control weeds, dense vegetative growth or weeds with a well-established root system.

### Preplant, At-Planting, Preemergence

**USE INSTRUCTIONS:** This product may be used to control broadleaf weeds and may be applied before, during or immediately after planting cotton with XtendFlex® technology. Refer to the "WEEDS CONTROLLED" section of the label booklet for XtendiMax™ With VaporGrip™ Technology for specific weeds controlled.

**RESTRICTIONS:** The maximum combined quantity of this product that may be applied for all preplant, at-planting, and preemergence applications is 44 fluid ounces (1.0 lb a.e. dicamba) per acre per season. The maximum application rate for a single, preplant, at-planting, or preemergence application must not exceed 44 fluid ounces (1.0 lb a.e. dicamba) per acre. Do not apply less than 22 fluid ounces (0.5 lb a.e. dicamba) per acre.

### Postemergence (In-crop)

**USE INSTRUCTIONS:** This product may be used to control broadleaf weeds in cotton with XtendFlex® technology. In-crop applications of this product can be made from emergence up to 7 days prior to harvest. The maximum and minimum rate for any single, in-crop application is 22 fluid ounces (0.5 lb a.e. dicamba) per acre. Using the appropriate application rate may reduce the selection for resistant weeds. For best performance, control weeds early when they are less than 4 inches. Monsanto Company does not warrant product performance of applications to labeled weeds greater than 4 inches in height. Sequential applications of this product may be necessary to control new flushes of weeds or on tough-to-control weeds. Allow at least 7 days between applications. A pre-harvest application of this product may be made up to 7 days before harvest.

Postemergence applications of this product mixed with adjuvants may cause a leaf response to cotton with XtendFlex® technology. The symptoms usually appear as necrotic spots on fully expanded leaves. To reduce the incidence and severity of the necrosis, consider increasing the spray volume to 15 GPA or greater and lower adjuvant rates. EC-based products that are tank mixed with products containing dicamba may increase the severity of the leaf damage.

**RESTRICTIONS:**

- The combined total applied from crop emergence up to 7 days prior to harvest must not exceed 88 fluid ounces (2.0 lb a.e. dicamba) per acre.
- The maximum single, in-crop application rate must not exceed 22 fluid ounces (0.5 lb a.e. dicamba).
- The combined total per year for all applications must not exceed 88 fluid ounces (2.0 lb a.e. dicamba) per acre. For example, if a preplant application of 44 fluid ounces (1.0 lb a.e. dicamba) per acre was made, then the combined total in-crop applications must not exceed 44 fluid ounces (1.0 lb a.e. dicamba) per acre.
- Allow at least 7 days between applications and allow at least 7 days between final application and harvest or feeding of cottonseed and cotton gin by-products.

**TANK-MIXING INSTRUCTIONS**

XtendiMax™ With VaporGrip™ Technology may only be tank-mixed with products that have been tested and found not to adversely affect the offsite movement potential of XtendiMax™ With VaporGrip™ Technology. A list of those products may be found at [www.xtendimaxapplicationrequirements.com](http://www.xtendimaxapplicationrequirements.com). DO NOT tank mix any product with XtendiMax™ With VaporGrip™ Technology unless:

- You check the list of tested products found not to adversely affect the offsite movement potential of XtendiMax™ With VaporGrip™ Technology at [www.xtendimaxapplicationrequirements.com](http://www.xtendimaxapplicationrequirements.com) no more than 7 days before applying XtendiMax™ With VaporGrip™ Technology; and
- The intended tank-mix product is identified on the list of tested products; and
- The intended products are not prohibited on either this supplemental label or the label of the tank mix product.
- Additional Warnings and Restrictions:
  - Some COC, HSOC and MSO adjuvants may cause a temporary crop response.
  - Do not tank mix products containing ammonium salts such as ammonium sulfate and urea ammonium nitrate.
  - Drift reduction agents (DRAs) can minimize the percentage of driftable fines. However, the applicator must check [www.xtendimaxapplicationrequirements.com](http://www.xtendimaxapplicationrequirements.com) to determine if the DRA is listed and check with the DRA manufacturer to determine if the DRAs will work effectively with the approved spray nozzle, spray pressure, and the desired spray solution.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MONSANTO MAKES NO RECOMMENDATION OR WARRANTY HEREIN REGARDING THE USE OF ANY PRODUCT THAT MAY APPEAR ON THE WEBSITE REFERENCED ABOVE, REGARDLESS OF WHETHER SUCH PRODUCT IS USED ALONE OR IN A TANK MIX WITH XTENDIMAX™ WITH VAPORGRIP™ TECHNOLOGY. BUYER AND ALL USERS ARE SOLELY RESPONSIBLE FOR ANY LACK OF PERFORMANCE, LOSS, OR DAMAGE IN CONNECTION WITH THE USE OR HANDLING OF ANY SUCH PRODUCT ALONE OR IN A TANK MIX WITH XTENDIMAX™ WITH VAPORGRIP™ TECHNOLOGY. See the section titled "LIMIT OF WARRANTY AND LIABILITY" herein for more information.

**WEED RESISTANCE MANAGEMENT**

Some naturally occurring weed biotypes that are tolerant (resistant) to dicamba may exist due to genetic variability in a weed population. Where resistant biotypes exist, the repeated use of herbicides with the same sites of action can lead to the selection for resistant weeds. Certain agronomic practices can delay or reduce the likelihood that resistant weed populations will develop and can be utilized to manage weed resistance once it occurs.

Do not use less than 22 fluid ounces per acre (0.5 lb a.e./A) of this product in a single application. Using the appropriate application rate can minimize the selection for resistant weeds.

Proactively implementing diversified weed control strategies to minimize selection for weed populations resistant to one or more herbicides is a best practice. A diversified weed management program may include the use of multiple herbicides with different sites of action and overlapping weed spectrum with or without tillage operations and/or other cultural practices. Research has demonstrated that using the labeled rate and directions for use is important to delay the selection for resistance.

The continued effectiveness of this product depends on the successful management of the weed resistance program; therefore, it is very important to perform the following actions.

To aid in the prevention of developing weeds resistant to this product, the following steps should be followed where practical:

- Scout fields before application to ensure herbicides and rates will be appropriate for the weed species and weed sizes present.
- Apply full rates of XtendiMax™ With VaporGrip™ Technology for the most difficult to control weed in the field at the specified time (correct weed size) to minimize weed escapes.
- Scout fields after application to detect weed escapes or shifts in weed species.
- Report any incidence of non-performance of this product against a particular weed species to your Monsanto retailer, representative or call 1-844-RRXTEND.
- If resistance is suspected, treat weed escapes with a herbicide having a site of action other than Group 4 and/or use non-chemical methods to remove escapes, as practical, with the goal of preventing further seed production.

Additionally, users should follow as many of the following herbicide resistance management practices as is practical:

- Use a broad spectrum soil-applied herbicide with other sites of action as a foundation in a weed control program.
- Utilize sequential applications of herbicides with alternative sites of action.
- Rotate the use of this product with non-Group 4 herbicides.
- Avoid making more than two applications of dicamba and any other Group 4 herbicides within a single growing season unless mixed with another mechanism of action with an overlapping spectrum for the difficult to control weeds.
- Incorporate non-chemical weed control practices, such as mechanical cultivation, crop rotation, cover crops and weed-free crop seeds, as part of an integrated weed control program.
- Thoroughly clean plant residues from equipment before leaving fields suspected to contain resistant weeds.

- Manage weeds in and around fields, during and after harvest to reduce weed seed production.

Contact the local agricultural extension service, Monsanto representative, agricultural retailer or crop consultant for further guidance on weed control practices as needed.

**APPLICATION EQUIPMENT AND TECHNIQUES**

DO NOT APPLY THIS PRODUCT TO COTTON WITH XTENDFLEX® TECHNOLOGY USING AERIAL SPRAY EQUIPMENT.

Apply this product using properly maintained and calibrated equipment capable of delivering the desired volumes.

**SPRAY DRIFT MANAGEMENT**

Do not allow herbicide solution to mist, drip, drift or splash onto desirable vegetation because severe injury or destruction to desirable broadleaf plants could result. The following drift management requirements must be followed.

**Controlling Droplet Size**

Off-target movement potential may be reduced by applying large droplets that provide sufficient coverage and control. Applying larger droplets can reduce drift potential, but will not prevent off-target movement if the application is made improperly, or under unfavorable environmental conditions (see the "Wind Speed and Direction", "Temperature and Humidity" and "Temperature Inversions" sections of this label).

- Nozzle type.** Use only Tee Jet® TT11004 nozzle with a maximum operating pressure of 63 psi when applying XtendiMax™ With VaporGrip™ Technology or any other approved nozzle found at [www.xtendimaxapplicationrequirements.com](http://www.xtendimaxapplicationrequirements.com). Do not use any other nozzle and pressure combination not specifically listed on this website.
- Hooded Sprayers.** Using a hooded sprayer in combination with approved nozzles may further reduce drift potential.
- Spray Volume.** Apply this product in a minimum of 10 gallons of spray solution per acre. Use a higher spray volume when treating dense vegetation. Higher spray volumes may also allow the use of larger nozzle orifices (sizes) which produce coarser spray droplets.
- Equipment Ground Speed.** Select a ground speed that will deliver the desired spray volume while maintaining the desired spray pressure, but do not exceed a ground speed of 15 miles per hour. Slower speeds generally result in better spray coverage and deposition on the target area.
- Spray boom Height.** Spray at the appropriate boom height based on nozzle selection and nozzle spacing, but do not exceed a boom height of 24 inches above target pest or crop canopy. Set boom to lowest effective height over the target pest or crop canopy based on equipment manufacturer's directions. Automated boom height controllers are recommended with large booms to better maintain optimum nozzle to canopy height. Excessive boom height will increase the potential for drift.

**Temperature and Humidity**

When making applications in low relative humidity or temperatures above 91 degrees Fahrenheit, set up equipment to produce larger droplets to compensate for evaporation. Larger droplets have a lower surface to volume ratio and can be impacted less by temperature and humidity. Droplet evaporation is most severe when conditions are both hot and dry.

**Temperature Inversions**

Do not apply this product during a temperature inversion. Drift potential can be high during a temperature inversion.

- During a temperature inversion, the atmosphere is very stable and vertical air mixing is restricted, which can cause small, suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions.
- Temperature inversions are characterized by increasing temperatures with altitude and are common on evenings and nights with limited cloud cover and light to no wind. Cooling of air at the earth's surface takes place and warmer air is trapped above it. They can begin to form as the sun sets and often continue into the morning.
- Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.
- The inversion will often dissipate with increased winds (above 3 MPH) or at sunrise when the surface air begins to warm (generally 3°F from morning low).

**Wind Speed and Direction**

- Drift potential is lowest between wind speeds of 3 to 10 miles per hour.
- Do not apply at wind speeds greater than 15 mph.
- For XtendiMax™ With VaporGrip™ Technology wind speed and direction restrictions see below table:

Wind speed	Application conditions and restrictions
<3 mph	Do not apply XtendiMax™ With VaporGrip™ Technology.
3-10 mph	Optimum application conditions for XtendiMax™ With VaporGrip™ Technology provided all other application requirements in this label are met.
>10 – 15 mph	Do not apply product when wind is blowing toward non-target sensitive crops.
> 15 mph	Do not apply XtendiMax™ With VaporGrip™ Technology.

**NOTE:** Local terrain can influence wind patterns. Every applicator must be familiar with local wind patterns and how they affect drift.

**PROTECTION OF SENSITIVE AREAS**

Maintain a 110 foot downwind buffer (when applying 22 fluid ounces of this product per acre) or a 220 foot downwind buffer (when applying 44 fluid ounces of this product per acre) between the last treated row and the closest downwind edge (in the direction in which the wind is blowing). If any of the following areas below are directly adjacent to the treated field, the areas listed below can be considered part of the buffer distance.

To maintain this required buffer zone:

- No application swath can be initiated in, or into an area that is within the applicable buffer distance.

The following areas may be included in the buffer distance calculation when adjacent to field edges:

- Roads, paved or gravel surfaces,
- Planted agricultural fields containing: corn, dicamba tolerant cotton, dicamba tolerant soybean, sorghum, proso millet, small grains and sugarcane. If the applicator intends to include such crops as dicamba tolerant cotton and/or dicamba tolerant soybeans in the buffer distance calculation, the applicator must confirm the crops are in fact dicamba tolerant and not conventional cotton and/or soybeans.
- Agricultural fields that have been prepared for planting.
- Areas covered by the footprint of a building, silo, or other man made structure with walls and or roof.

#### Non-target Susceptible Crops

**Failure to follow the requirements in this label could result in severe injury or destruction to desirable sensitive broadleaf crops and trees when contacting their roots, stems or foliage.**

- Do not apply under circumstances where off-target movement may occur to food, forage, or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.
- Do not allow contact of herbicide with foliage, green stems, exposed non-woody roots of crops, and desirable plants, including beans, cotton, flowers, fruit trees, grapes, ornamentals, peas, potato, soybean, sunflower, tobacco, tomato, and other broadleaf plants because severe injury or destruction may result, including plants in a greenhouse.
- Small amounts of dicamba that may not be visible may injure susceptible broadleaf plants.
- Applicators are required to ensure that they are aware of the proximity to non-target susceptible crops, and to avoid potential adverse effects from drift of XtendiMax™ With VaporGrip™ Technology.

**Before making an application, the applicator must survey the application site for neighboring non-target susceptible crops. The applicator must also consult sensitive crop registries to identify any commercial specialty or certified organic crops that may be located near the application site.**

**DO NOT APPLY this product when the wind is blowing toward adjacent commercially grown dicamba sensitive crops, including but not limited to, commercially grown tomatoes and other fruiting vegetables (EPA crop group 8), cucurbits (EPA crop group 9), and grapes.**

#### Application Awareness

AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR

The interaction of equipment and weather related factors must be monitored to maximize performance and on-target spray deposition. The applicator is responsible for considering all of these factors when making a spray decision. The applicator is responsible for compliance with state and local pesticide regulations, including any state or local pesticide drift regulations.

#### Proper spray system equipment cleanout

Minute quantities of dicamba may cause injury to non-dicamba-tolerant soybeans and other sensitive crops (see the "Non-target Susceptible Crops" section of this label for more information).

Clean equipment immediately after using this product, using a triple rinse procedure as follows:

1. After spraying, drain the sprayer (including boom and lines) immediately. Do not allow the spray solution to remain in the spray boom lines overnight prior to flushing.
2. Flush tank, hoses, boom and nozzles with clean water.
3. Inspect and clean all strainers, screens and filters.
4. Prepare a cleaning solution with a commercial detergent or sprayer cleaner or ammonia according to the manufacturer's directions.
5. Take care to wash all parts of the tank, including the inside top surface. Start agitation in the sprayer and thoroughly recirculate the cleaning solution for at least 15 minutes. All visible deposits must be removed from the spraying system.
6. Flush hoses, spray lines and nozzles for at least 1 minute with the cleaning solution.
7. Repeat above steps for two additional times to accomplish an effective triple rinse.
8. Remove nozzles, screens and strainers and clean separately in the cleaning solution after completing the above procedures.
9. Appropriately dispose of rinsate from steps 1-7 in compliance with all applicable laws and regulations.
10. Drain sump, filter and lines.
11. Rinse the complete spraying system with clean water.

All rinse water must be disposed of in compliance with local, state, and federal requirements.

#### CROP ROTATIONAL RESTRICTIONS

No rotational cropping restrictions apply when rotating to Roundup Ready® 2 Xtend™ Soybeans or cotton seed with XtendFlex® technology (including Bollgard® 3 XtendFlex® Cotton, Bollgard II® XtendFlex® Cotton, or XtendFlex® Cotton). For other crops the interval between application and planting rotational crop is given below. When counting days from the application of this product, do not count days when the ground is frozen. Planting at intervals less than specified below may result in crop injury. Moisture is essential for the degradation of this herbicide in soil. If dry weather prevails, use cultivation to allow herbicide contact with moist soil.

#### Planting/replanting restrictions for XtendiMax™ With VaporGrip™ Technology applications of 33 fluid ounces per acre or less

For corn, cotton (except cotton seed with XtendFlex® technology), sorghum, and soybean (except Roundup Ready® 2 Xtend™ Soybean), follow the planting restrictions in the directions for use for preplant application in **Section 10. Crop-Specific Information** of the label booklet. Do not plant barley, oat, wheat, and other grass seedlings for 15 days for every 11 fluid ounces of this product applied per acre east of the Mississippi River and 22 days for every 11 fluid ounces per acre applied west of the Mississippi River. No planting restrictions apply beyond 120 days after application of this product.

#### Planting/replanting restrictions for applications of more than 33 fluid ounces and up to 44 fluid ounces of XtendiMax™ With VaporGrip™ Technology per acre

Wait a minimum of 120 days after application of this product before planting corn, sorghum and cotton (except cotton seed with XtendFlex® technology) east of the Rocky Mountains and before planting all other crops (except Roundup Ready® 2 Xtend™ Soybean) grown in areas receiving 30 inches or more rainfall annually. Wait a minimum of 180 days before planting crops in areas with less than 30 inches of annual rainfall. Wait a minimum of 30 days for every 22 fluid ounces of this product applied per acre before planting barley, oat, wheat, and other grass seedlings east of the Mississippi River and 45 days for every 22 fluid ounces of this product applied per acre west of the Mississippi River.

#### LIMIT OF WARRANTY AND LIABILITY

Monsanto Company ("Company") warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in this supplemental label ("Directions") when used in accordance with the Directions under the conditions described therein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein. Specifically, and without limiting the foregoing, MONSANTO MAKES NO RECOMMENDATION OR WARRANTY HEREIN REGARDING THE USE OF ANY PRODUCTS THAT MAY APPEAR ON THE WEBSITE REFERENCED IN THE TANK-MIXING INSTRUCTIONS HEREIN, REGARDLESS OF WHETHER SUCH PRODUCT IS USED ALONE OR IN A TANK MIX WITH XTENDIMAX™ WITH VAPORGRIP™ TECHNOLOGY. BUYER AND ALL USERS ARE SOLELY RESPONSIBLE FOR ANY LACK OF PERFORMANCE, LOSS, OR DAMAGE IN CONNECTION WITH THE USE OR HANDLING OF ANY SUCH PRODUCT ALONE OR IN A TANK MIX WITH XTENDIMAX™ WITH VAPORGRIP™ TECHNOLOGY.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, tort, or otherwise.

To the extent consistent with applicable law, buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those expressly recommended by Company in the Directions, application to or contact with desirable vegetation, failure of this product to control weed biotypes which develop resistance to dicamba, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those expressly recommended by Company in the Directions in or on the soil, crop or treated vegetation.

This Company does not warrant any product reformulated or repackaged from this product except in accordance with this Company's stewardship requirements and with express written permission from this Company.

For in-crop (over-the-top) uses on crops within the Roundup Ready Xtend® Crop System, crop safety and weed control performance are not warranted by Company when this product is used in conjunction with "brown bag" or "bin run" seed saved from previous year's production and replanted.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Upon opening and using this product, buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement.

These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

©2016

MONSANTO COMPANY  
800 N. LINDBERGH BLVD.  
ST. LOUIS, MISSOURI 63167 USA

11/15/2016

35008P3-9